

Calcium SC Plus™

A calcium suspension concentrate for foliar application with added zinc and boron

FERTILIZER GROUP 2

ACTIVE INGREDIENTS

Calcium (Ca) 242 g/kg (400 g/ℓ)

Zinc (Zn) 18 g/kg (30 g/ℓ)

Boron (B) 2 g/kg (3 g/ℓ)

SG = 1.65 @ 20°C

Contents

Batch number

Manufacture date

Contents

Batch number

Manufacture date

Sineria

SINERIA HOLLAND BV
Randwycksingel 20-A015
6229 EE Maastricht The Netherlands
www.sineria.com

- **Calcium SC Plus™** should preferably be applied in the early morning or late afternoon.
- It is advisable to do a miscibility test prior to mixing **Calcium SC Plus™** with other chemicals.
- Consult with your chemical distributor should crop specific programmes or any other information be required regarding the use of **Calcium SC Plus™**.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAXIMUM RATE PER HA.	RATE PER 100ℓ WATER	REMARKS
Apples	5 ℓ	200 - 250 mℓ	Apply from petal fall to one month before harvest with 10 - 14 day intervals. Apply as a full cover spray in 1 000 - 3 000 ℓ water/ha. Avoid run-off.
Avocados and Mangoes	5 ℓ	250 mℓ	Apply from fruit set to one month before harvest, with 10 - 14 day intervals. Apply in 2 000 ℓ water/ha.
Bananas	2 ℓ	—	Apply every 10 - 14 days to one month before harvest. Apply in 30 ℓ water/ha.
Citrus	5 ℓ	125 mℓ	Apply from fruit set. Do not apply within one month before harvest. Apply in 4 000 ℓ water/ha.
Cucurbits (Cucumbers, Melons)	2 - 4 ℓ	—	Apply with 7 - 10 day intervals from first fruit set to one month before harvest. Repeat if necessary. Apply in 200 - 400 ℓ water/ha.
Deciduous Fruits	5 ℓ	250 mℓ	Apply with 7 - 14 day intervals from petal fall to one month before harvest. Apply in 2 000 ℓ water/ha. If conditions are conducive to fruit russet (variety, weather conditions, etc) start applications 6 weeks after petal fall.
Grapes	5 ℓ	—	Apply every 10 - 14 days from fruit set in 1 000 ℓ water/ha.
Groundnuts	3 - 5 ℓ	—	Start application at 4 - 6 leaf stage. If necessary, repeat application with 10 - 14 day intervals. Apply in 30 - 200 ℓ water/ha.
Nuts (Macadamias, Pecans)	5 ℓ	250 mℓ	Start application 7 days after petal fall. Repeat if necessary with 10 - 14 day intervals. Do not apply within one month before harvest. Apply in 1 000 - 2 000 ℓ water/ha.
Peppers (field grown)	5 ℓ	—	Apply from 4 to 6 leaf stage to one month before harvest. Repeat applications every 7 - 10 days. Apply in 50 - 500 ℓ water/ha. Start
Strawberries (field grown)	3 - 5 ℓ	—	application early in the new season until one month before harvest. Apply 3 applications with 10 - 14 day intervals in 300 - 500 ℓ water/ha.
Tomatoes (field grown)	5 ℓ	—	Start application at the end of flowering on second truss until one month before harvest. If necessary repeat with 7 - 10 day intervals. Apply a maximum of 5 applications per season. Apply in 50 - 500 ℓ water/ha.
Vegetables (field grown) (Cabbage, Carrots, Lettuce, etc.)	3 - 5 ℓ	—	Apply from 4 - 9 leaf stage until one month before harvest. Repeat with 10 - 14 day intervals. Apply in 50 - 500 ℓ water/ha.