

COMBI-PRO™

COMBI-PRO™ is an organic fertilizer for use as a foliar feed to maintain or correct zinc, iron, copper and manganese levels in plants

FERTILIZER GROUP 2

ACTIVE INGREDIENTS

Zinc	(Zn)	20,9 g/kg	(27,1 g/l)
Iron	(Fe)	26,6 g/kg	(34,6 g/l)
Copper	(Cu)	10,0 g/kg	(13,0 g/l)
Manganese	(Mn)	17,3 g/kg	(22,5 g/l)
Boron	(B)	3,1 g/kg	(4,0 g/l)
Molybdenum	(Mo)	0,12 g/kg	(0,15 g/l)

SG = 1,27 ± 0,02

L-amino acid chelated

Sineria

SINERIA HOLLAND BV
Randwycksingel 20-A015
6229 EE Maastricht The Netherlands
www.sineria.com

Contents • Batch number • Manufacture date

Blank area for labeling information.

COMBI-PRO™ is compatible with most agricultural remedies. However, do **NOT** apply **COMBI-PRO™**:

- in combination with copper containing fungicides, phosphates or phosphonates
- within 30 days after application of any copper fungicides
- in combination with highly alkaline material such as lime sulphur or certain boron products.

COMBI-PRO™ should preferably be applied in the early morning or late afternoon. Do not apply to plants that are undergoing a period of moisture or heat stress.

For optimum results, **COMBI-PRO™** should be used in a programme and in combination with other balanced fertilizers and foliar feeds. Consult your chemical distributor should crop specific programmes or any other information be required about the use of **COMBI-PRO™**.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAX. RATE PER HA.	RATE PER 100ℓ WATER	REMARKS
Apples and Pears	3 ℓ	100 mℓ	Apply as a full cover spray in 1 000 to 3 000 ℓ spray water in spring when the first leaves appear. Repeat after harvest.
Avocado's	5 ℓ	150 mℓ	Start application in the spring, and repeat 2 to 3 times if necessary. Apply as a full cover spray in 1 000 to 3 000 ℓ spray water per hectare.
Cruciferae, Lettuce, Beetroot, Celery	1,5 ℓ		Apply 2 to 3 sprays, starting 4 weeks after transplant. Apply in 300 to 500 ℓ water per hectare.
Cucurbits (Pumpkins, Cucumbers, etc.)	3 ℓ		Start application directly after fruit set. Repeat every 7 to 10 days if necessary. Apply in 500 ℓ water per hectare.
Citrus	5 ℓ	150 mℓ	Start application early in spring. Repeat every 7 to 10 days (2 to 3 applications). Apply in 1 000 to 3 000 ℓ water per hectare.
Cotton	3 ℓ		Start application when plants reach a height of 30 cm. Repeat 2 to 3 times. Apply in 500 ℓ water per hectare.
Grapes	2 - 5 ℓ		Start application directly after fruit set. Apply 3 to 4 sprays every 10 days. Apply in 1 000 to 1 500 ℓ water per hectare.
Litchi and Mangoes	5 ℓ	150 mℓ	Start application shortly before flowering. Apply 2 to 3 sprays at 10-day intervals. Apply in 1 000 to 3 000 ℓ water per hectare.
Potatoes	3 ℓ		Start application at tuber initiation. Apply 3 to 4 sprays at 10 to 14 day intervals. Apply in 500 ℓ water per hectare.
Soybeans, Groundnuts and Green Beans	1,5 ℓ		Start application at flowering. Apply 2 sprays at 7-day intervals. Apply in 300 ℓ water per hectare.
Stone Fruit (Plums, Peaches, Nectarines)	3 - 5 ℓ	150 mℓ	Start application directly after fruit set. Apply 2 to 3 sprays at 10-day intervals. Apply in 1 000 to 3 000 ℓ water per hectare.
Sugarcane	3 ℓ		Start application in early spring. Apply 2 sprays at 10-day intervals. Apply in 500 to 1 500 ℓ water per hectare.
Tomatoes and Peppers	3 ℓ		Start application at first set. Repeat at 7 to 10 day intervals. Apply in 500 ℓ water per hectare.
Other Crops	3 ℓ		Apply as a medium cover spray and repeat at 14-day intervals. Apply in 500 ℓ water per hectare.

Use lower rates as a maintenance spray and higher rates to correct deficiencies.