

COP-PRO™

COP-PRO™ is an organic copper fertilizer for use as a foliar feed to maintain or correct copper levels in plants

FERTILIZER GROUP 2

ACTIVE INGREDIENT

Copper (Cu) 80 g/kg (106 g/ℓ)

SG = 1,32 ± 0,02

L-amino acid chelated

Sineria

SINERIA HOLLAND BV
Randwycksingel 20-A015
6229 EE Maastricht The Netherlands
www.sineria.com

Contents • Batch number • Manufacture date

--

COP-PRO™ is compatible with most agricultural remedies. However, do **NOT** apply **COP-PRO™**:

- in combination with copper containing fungicides, phosphates or phosphonates
- within 30 days after application of any copper fungicides
- in combination with highly alkaline material such as lime sulphur or certain boron products.

COP-PRO™ should preferably be applied in the early morning or late afternoon. Do not apply to plants that are undergoing a period of moisture or heat stress.

For optimum results, **COP-PRO™** should be used in a programme and in combination with other balanced fertilizers and foliar feeds. Consult your chemical distributor should crop specific programmes or any other information be required about the use of **COP-PRO™**.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAX. RATE PER HA.	RATE PER 100ℓ WATER	REMARKS
Citrus	750 mℓ	20 - 25 mℓ	Apply in September, before flowering. Repeat 3 to 4 weeks later if necessary. Apply in a minimum of 1000 to 3000 ℓ water per hectare.
Onions	500 mℓ	75 - 100 mℓ	Apply from golf ball size bulbs, and repeat 14 days later. Apply in a minimum of 500 ℓ water per hectare.
Pastures	250 - 500 mℓ		Apply early in the season before spring growth starts. Repeat between cuts of silage or hay. Important: Keep all livestock out of sprayed pasture for two weeks after application.
Pome, Stone, Deciduous and Sub-tropical Fruit	750 mℓ	20 - 25 mℓ	Apply with bud swell and repeat application at 75% petal fall. Apply in a minimum of 1000 to 3000 ℓ water per hectare.
Sugarbeet	200 mℓ	75 - 100 mℓ	For best results, apply early in the season when there is sufficient leaves to absorb the spray. Apply 1 to 2 applications in a minimum of 200 ℓ water per hectare.
Wheat, Barley, Oats, Maize	200 mℓ	75 - 100 mℓ	Apply 1 to 2 applications early in the season when there is enough leaves to absorb the spray. Apply in a minimum of 200 ℓ water per hectare.

Use lower rates as a maintenance spray and higher rates to correct deficiencies.