

# Ferro Col Copper 15%<sup>TM</sup>

A water-soluble powder copper EDTA chelate  
for use as a soil application or foliar spray

## FERTILIZER GROUP 2

### ACTIVE INGREDIENT

Copper (Cu) 150 g/kg

*EDTA Chelated*

**Sineria**

SINERIA HOLLAND BV  
Randwycksingel 20-A015  
6229 EE Maastricht The Netherlands  
[www.sineria.com](http://www.sineria.com)

Contents • Batch number • Manufacture date

Blank area for labeling information.

**FOLIAR APPLICATION:** Apply in sufficient water to obtain adequate coverage of the foliage. Apply early morning or late afternoon for the best results.

**SOIL APPLICATION:** Apply in sufficient water to obtain an even distribution. Incorporate into the soil immediately after application.

**COMPATIBILITY:** **Ferro Col Copper 15%™** is compatible with most agricultural chemicals.

**WARNING:** The use of a wetter or buffer is not recommended as phytotoxicity may occur.

**Consult with a qualified agronomist or agricultural advisor on the necessity of using it, before applying this fertilizer.**

## **DIRECTIONS FOR USE: USE ONLY AS DIRECTED**

<b>CROP</b>	<b>DOSAGE PER HECTARE</b>	<b>REMARKS</b>
<b>Citrus</b> and other fruit crops	Foliar: 0,5 kg Soil:	Apply early in the season and repeat 3–4 weeks later. Avoid spraying during bloom and just before harvest. Apply in a minimum of 500 ℓ water per hectare.
<b>Wheat, Barley, Oats, Maize and Onions</b>	0,5 - 1 kg	Apply immediately pre-planting in a minimum of 200 ℓ water per hectare. Apply annually until the deficiency is corrected. Thereafter apply every three years to prevent recurrence.
	Foliar:0,25 - 0,5 kg	If possible apply before deficiencies occur and repeat 3–4 weeks later if necessary. Apply early in the growing season, as soon as the crop has sufficient leaves to absorb the spray. Apply in a minimum of 200 ℓ water per hectare.
<b>Conifers</b>	Foliar:1 kg	Apply by air, ULV or conventional sprayer to second generation trees. Apply in 200–500 ℓ water per hectare depending on tree size.
<b>Pasture</b>	Re-seeding: 4 kg	Spray and incorporate into the soil prior to re-seeding.
	Established pasture: 0,5 - 1 kg	Spray early in the season before spring growth starts, and again between cuts of silage or hay. <b>Keep sheep out of recently sprayed pasture.</b>