

Ferro Col Manganese 13%TM

A water-soluble powder manganese EDTA chelate
for use as a foliar spray

FERTILIZER GROUP 2

ACTIVE INGREDIENT

Manganese (Mn) 130 g/kg

EDTA Chelated

Sineria

SINERIA HOLLAND BV
Randwycksingel 20-A015
6229 EE Maastricht The Netherlands
www.sineria.com

Contents • Batch number • Manufacture date

--

FOLIAR APPLICATION: Apply in sufficient water to obtain adequate coverage of the foliage. Apply early morning or late afternoon for the best results. Manganese deficiency is most likely to occur in highly leached, very sandy, alkaline, calcareous or heavy limed soils. Manganese deficiency is seldom encountered in loam or heavy acid soils.

COMPATIBILITY: **Ferro Col Manganese 13%™** is compatible with most foliar applied agrochemicals.

Consult with a qualified agronomist or agricultural advisor on the necessity of using it, before applying this fertilizer.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	DOSAGE PER HECTARE	REMARKS
Stone Fruit: Apricots, Peaches, Plums	1 kg	Apply as a foliar spray after harvest. Apply in a minimum of 500 ℓ water per hectare.
Pome Fruit: Apples, Pears	1 kg	Apply as a foliar spray before blossom and repeat after harvest. Apply in a minimum of 500 ℓ water per hectare.
Citrus	1 kg	Apply as a foliar spray to new growth flush in early spring. Repeat 3–4 weeks later. Apply in a minimum of 500 ℓ water per hectare.
Field Crops: Maize, Beans, Peas, Soya beans, Groundnuts, Wheat, Barley, Oats, Cruciferae, Onions, Tomatoes	1,2 kg	Apply early in the growing season when the crop has sufficient leaves to absorb the spray. Repeat 3–4 weeks later if necessary. Apply in a minimum of 200 ℓ water per hectare.
Olives	1,2 kg	Apply during late spring as a foliar spray in a minimum of 500 ℓ water per hectare.
Pastures	4 kg	Apply early spring and repeat annually. Apply in a minimum of 200 ℓ water per hectare.