

Ferro Col Zinc 15%TM

A water-soluble powder zinc EDTA chelate
for use as a soil application or foliar spray

FERTILIZER GROUP 2

ACTIVE INGREDIENT

Zinc (Zn) 150 g/kg

EDTA Chelated

Sineria

SINERIA HOLLAND BV
Randwycksingel 20-A015
6229 EE Maastricht The Netherlands
www.sineria.com

Contents • Batch number • Manufacture date

Blank area for product information.

FOLIAR APPLICATION: Apply in sufficient water to obtain adequate coverage of the foliage. Apply early morning or late afternoon for the best results.

SOIL APPLICATION: Apply in sufficient water to obtain an even distribution. Incorporate into the soil immediately after application.

COMPATIBILITY: **Ferro Col Zinc 15%™** is compatible with the majority of commonly used pesticides and fertilizers.

Consult with a qualified agronomist or agricultural advisor on the necessity of using it, before applying this fertilizer.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	DOSAGE PER HECTARE	REMARKS
Field Crops: Maize, Beans, Peas, Soyabeans, Groundnuts, Wheat, Barley, Oats, Cruciferae, Onions, Tomatoes	Soil: 2-4 kg	Apply immediately pre-planting. Apply annually until the deficiency is corrected, and then every 3 years to prevent recurrence.
	Foliar: 1-2 kg	Apply early in the growing season, as soon as the crop has sufficient leaves to absorb the spray. If possible apply before deficiencies occur. Apply in a minimum of 200 ℓ water/ha.
Pecans	Foliar: 0,5 kg	Apply as a full cover spray at full leaf in the spring. Repeat twice with 14 day intervals. Apply in a minimum of 500 ℓ water/ha.
Citrus	Foliar: 1-2 kg	Spray the new flush of growth in spring. Repeat when necessary but avoid spraying during and prior to harvest. Apply in a minimum of 500 ℓ water/ha.
Pome Fruit: Apples and Pears	Foliar: 1 kg	Spray before blossom and repeat after harvest. Apply in a minimum of 500 ℓ water/ha.
Stone Fruit: Apricots, Peaches, Plums	Foliar: 1 kg	Spray after harvest. Do not exceed the recommended rate or leaf scorch may occur. Apply in a minimum of 500 ℓ water/ha.
	Soil: 0,5-1 kg/tree	Apply as a drench to the drip zone area in case of severe deficiencies. Dosage rate depends on tree size.
Olives	Foliar: 1 kg	Spray in spring when new growth is starting, and repeat 3 - 4 weeks later. Apply in a minimum of 500 ℓ water/ha.
Cherries	Foliar: 0,5-1 kg	Apply as a full cover spray just after bloom. Repeat twice with 2 - 3 week intervals. Apply in a minimum of 500 ℓ water/ha.