

GLUCO-Mag™

GLUCO-Mag™ is an organic fertilizer for use as a foliar feed to maintain or correct magnesium levels in plants

ACTIVE INGREDIENTS

Magnesium (Mg)	50 g/kg	(65 g/l)
Boron (B)	3 g/kg	(4 g/l)
L-amino acids	20 g/kg	(26 g/l)

SG = 1,28 ± 0,02

Gluconate Chelated

Sineria

SINERIA HOLLAND BV
Randwycksingel 20-A015
6229 EE Maastricht The Netherlands
www.sineria.com

Contents • Batch number • Manufacture date

Blank area for labeling information.

For optimum results, **GLUCO-Mag™** should be used in a programme and in combination with other balanced fertilizers and foliar feeds. **GLUCO-Mag™** is compatible with most agricultural remedies. It is however advisable to do a miscibility test prior to mixing with other chemicals. **Do not mix GLUCO-Mag™** with highly alkaline material such as Lime Sulphur and Bordeaux mixture, or with any phosphate-containing fertilizers. **GLUCO-Mag™** should preferably be applied in the early morning or late afternoon. Do not apply to plants undergoing a period of moisture or heat stress. Consult your chemical distributor should crop specific programmes or any other information be required about the use of **GLUCO-Mag™**.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAX. RATE PER HA.	RATE PER 100ℓ WATER	REMARKS
Citrus	8 ℓ	300–600 mℓ	1 to 2 applications as a high volume spray in a minimum of 1000 ℓ water per hectare.
Cotton	3,5 ℓ	300–600 mℓ	1 to 2 applications. Start application as soon as deficiency symptoms appear. Apply in 300–500 ℓ water per ha.
Grapes	6 ℓ	300–600 mℓ	Start application two weeks after flowering. Repeat with 14 day intervals in a minimum of 1000 ℓ water per ha.
Grains, Potatoes, Peas, Canola	3,5 – 4 ℓ	300–600 mℓ	Start application when deficiency symptoms appear. Repeat if necessary. Apply in a minimum of 300–500 ℓ water per hectare.
Onions	4 ℓ	300–600 mℓ	3 to 4 applications. Apply in a minimum of 500 ℓ water per hectare.
Tea	7,5 ℓ	300–600 mℓ	Apply as a ground spray as soon as deficiency symptoms appears in 200–500 ℓ water per hectare. Aerial application in 30 - 45 ℓ water per hectare.
Tobacco	3,5 ℓ	300–600 mℓ	3 to 4 applications at 10 to 14 day intervals. Apply in 500 ℓ water per hectare.
Tomatoes, Peppers, Cabbage, Cucurbits, Eggfruit, Lettuce	4 – 6 ℓ	300–600 mℓ	3 to 4 applications. Start application when deficiency symptoms appear. Apply in a minimum of 200–400 ℓ water per hectare.
Other Crops	1,5 – 5 ℓ	300–600 mℓ	Start application when deficiency symptoms appear and repeat if necessary. Apply in 300–500 ℓ water per hectare.

Use lower rates as a maintenance spray and higher rates to correct deficiencies.