

MULTI-TRACE™



5ℓ

MULTI-TRACE™

is a fertilizer for use as a foliar feed to maintain or correct zinc, iron, copper and manganese levels in plants. Also contains boron and molybdenum.

Active ingredients:

Zinc (Zn) 33 g/kg (43 g/ℓ)

Iron (Fe) 13 g/kg (16 g/ℓ)

Copper (Cu) 9 g/kg (12 g/ℓ)

Manganese (Mn) 16 g/kg (21 g/ℓ)

Boron (B) 0,3 g/kg (0,4 g/ℓ)

Molybdenum (Mo) 0,08 g/kg (0,1 g/ℓ)

Aminogluconate Chelated

SG (20 °C) = 1,30

Fertilizer Group 2

Sineria

SINERIA HOLLAND BV
Randwycksingel 20-A015
6229 EE Maastricht The Netherlands
www.sineria.com

MULTI-TRACE™ is compatible with most agricultural remedies. However, do **NOT** apply **MULTI-TRACE™**:

- in combination with copper containing fungicides, phosphates or phosphonates
- within 30 days after application of any copper fungicides
- in combination with highly alkaline material such as lime sulphur or certain boron products.

MULTI-TRACE™ should preferably be applied in the early morning or late afternoon. **Do not apply to plants that are undergoing a period of moisture or heat stress.**

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

For optimum results **MULTI-TRACE™** should be used in a programme and in combination with other balanced fertilizers and foliar feeds. Use lower rates as a maintenance spray and higher rates to correct deficiencies.

Consult your chemical distributor should crop specific programmes or any other information be required regarding the use of **MULTI-TRACE™**.

CROP	MAXIMUM RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
Cereals and Canola	2,5 ℓ	500–600 mℓ	Start application at 3–4 leaf stage and repeat before flowering. Apply in 300 ℓ water per hectare.
Avocados	5 ℓ	150–165 mℓ	Start application in the spring, and repeat 2–3 times with 7–14 day intervals. Apply as a full cover spray in 1 000 – 3 000 ℓ water per hectare.
Cruciferae, Lettuce, Beetroot, Celery	1,5 ℓ	200–300 mℓ	Apply 2–3 sprays after transplant, as soon as plants are established. Apply in 300 – 500 ℓ water per hectare.
Cucurbits	3 ℓ	500–600 mℓ	Start application directly after fruit set. Repeat every 7–10 days if necessary. Apply in 500 ℓ water per hectare.
Citrus	5 ℓ	150 mℓ	Start application early in spring. Repeat 2–3 times every 7–10 days. Apply in 1 000 – 3 000 ℓ water per hectare.
Cotton	3 ℓ	600 mℓ	Start application when plants are 30 cm high. Repeat 2–3 times in 500 ℓ water per hectare.
Grapes and Persimmons	5 ℓ	150–250 mℓ	Start application at 10 cm shoot growth. Apply 3–4 sprays every 10 days in 1 000 – 1 500 ℓ water per hectare.
Litchi and Mangoes	5 ℓ	150–250 mℓ	Start application shortly before flowering. Apply 2–3 sprays at 10 day intervals in 1 000 – 3 000 ℓ water per hectare.
Potatoes	3 ℓ	500–600 mℓ	Start application at tuber initiation. Apply 3–4 sprays at 10–14 day intervals. Apply in 500 ℓ water per hectare.
Soybeans, Groundnuts and Green Beans	1,5 ℓ	300–500 mℓ	Start application at flowering. Apply 2 sprays at 7 day intervals. Apply in 300 ℓ water per hectare.
Pome and Stone Fruit	3 ℓ	100 mℓ	Start application directly after fruit set. Apply 2–3 at 10 day intervals. Apply in 1 000 – 3 000 ℓ water per hectare.
Sugarcane	3 ℓ	200–500 mℓ	Start application in early spring. Apply 2 sprays at 10 day intervals. Apply in 500 - 1 500 ℓ water per hectare.
Tomatoes and Peppers	3 ℓ	500–600 mℓ	Start application at first set. Repeat at 7–10 day intervals. Apply in 500 ℓ water per hectare.