

DIRECTIONS FOR USE: USE ONLY AS RECOMMENDED

SINERIA Calmag +TE™ is recommended for use during plant growth periods that require relatively high Calcium and Magnesium levels and during periods of stress when the uptake and movement of calcium in the plant is reduced.

SINERIA Calmag +TE™ is specifically formulated to correct cation imbalances that may occur in plants.

SINERIA Calmag +TE™ is a completely water soluble fertilizer and chlorine free.

SINERIA Calmag +TE™ is readily taken up by the plant and is formulated as both a foliar feed and for fertigation.

SINERIA Calmag +TE™ contains amino acids.

For optimum results, **SINERIA Calmag +TE™** should be used in a program in combination with the other **SINERIA** range of fertilizers.

SINERIA Calmag +TE™ is compatible with most agricultural remedies. It is however advisable to do a miscibility test prior to mixing with other chemicals. **Do not apply with products containing high levels of phosphorus.**

The rate of application is dependent on the soil fertility and reaction required. The lower rates should be used for maintenance, while the higher rates will correct nutritional deficiencies.

SINERIA Calmag +TE™ should preferably be applied in the early morning or late afternoon. Do not apply to plants that are undergoing a period of moisture or heat stress.

Consult your chemical distributor, consultant or manufacturer should crop specific programs or any other information be required around the use of the **SINERIA** range of fertilizers.

APPLICATION	RATE PER HECTARE	RATE PER 100 ℓ WATER	REMARKS
Field and Vegetable crops: Beans, Brassicae, Canola, Cereals, Celery, Cotton, Cucurbits, Garlic, Groundnuts, Lettuce, Maize, Onions, Peas, Peppers, Potatoes, Soybeans, Sweetcorn, Tobacco, Tomatoes, Wheat	1 - 5 ℓ	1 - 2 ℓ	Apply as a full cover spray directly onto fruit. Apply at 1 to 2 week intervals.
	1 - 5 ℓ	1 - 3 ℓ per 30 ℓ water	Aerial application: Apply at 1 to 2 week intervals as required.
Ornamental crops	1 - 5 ℓ	1 - 2 ℓ	Apply every 7 to 14 days as a full cover spray.
Large trees, Vines and Orchards: Apples, Apricots, Avocados, Bananas, Coffee, Cherries, Citrus, Grapes, Guavas, Litchis, Macadamias, Mangoes, Olives, Peaches, Pears, Pineapples, Plums	4 - 6 ℓ	1 - 2 ℓ	Apply at 1 to 2 week intervals. In case of deficiencies apply directly onto the fruit.
	Fertigation	5 - 10 ℓ	250 ml - 2 ℓ