

**SIVAMEX PLUS** fluke and worm drench for cattle, sheep, and goats is a yellow suspension.

#### COMPOSITION

Contains per ml:

Levamisole Hcl BP Vet, 1.5% w/v

Oxyclozanide BP Vet, 3% w/v

Cobalt Sulphate 0.382% w/v

#### INDICATIONS

**SIVAMEX 1.5%** is used in the treatment and control of gastrointestinal worms and lungworms in cattle, sheep, goats and pigs.

It is safe in animals of all ages and can be used in pregnant and lactating animals.

#### CONTRA-INDICATIONS, WARNING

1. Due care should be exercised in pregnant animals.

It is preferable to administer the product in ewes 6 weeks before lambing.

2. If animals are dosed during lactation, do not use milk for human consumption for 72 hrs after the last dose.

3. Do not slaughter stock for human consumption for 7 days after the last dose.

#### Meat and Milk Withdrawal

Do not administer this product to sheep and goats producing milk for human consumption.

Milk from treated cows should only be consumed 72 hrs from the last treatment.

Animals should only be slaughtered for human consumption 28 days from the last treatment.



**Combined Fluke and roundworms drench for Cattle, Sheep and Goats**

**Contains Cobalt for improved growth and performance  
Removes bowel roundworm segments completely**

#### DOSAGE AND ADMINISTRATION

Shake well before use.

SIVAMEX 1.5% should be administered as an oral drench at a dose rate of 7.5 mg per Kg body weight using a dosing gun or any other appropriate measure.

#### Dosage

##### Cattle

50 kg: 25 ml

100 kg: 50 ml

150 kg: 75 ml

200 kg: 100 ml

250 kg: 125 ml

300 Kg: 150 ml

##### Sheep/Goats

10 kg: 5 ml

20 kg: 10 ml

30 kg: 15 ml

40 kg: 20 ml

50 kg: 25 ml

60 kg: 30 ml

Batch No.:

Mfg. Date:

Exp. Date:

#### CAUTION

For animal treatment only.

Keep out of reach of children.

Store in a dark place below 30°C

#### Manufactured For:

Sineria East Africa Ltd  
P.O. Box , Nairobi, Kenya.

#### Manufactured by:

Stedam Pharma Manufacturing Ltd  
P.O. Box 35240-00200, Nairobi, Kenya