

CALMAG-PRO™

CALMAG-PRO is an organic fertilizer for use as a foliar feed to maintain or correct calcium and magnesium levels in plants

FERTILIZER GROUP 1

ACTIVE INGREDIENTS

Calcium (Ca)	48,1 g/kg	(62,6 g/l)
Magnesium (Mg)	16,1 g/kg	(20,9 g/l)

SG = 1,32 ± 0,02

L-amino acid chelated

Sineria

SINERIA HOLLAND BV
Randwycksingel 20-A015
6229 EE Maastricht The Netherlands
www.sineria.com

Contents • Batch number • Manufacture date

Blank area for labeling information.

CALMAG-PRO™ is compatible with most agricultural remedies. However, do **NOT** apply **CALMAG-PRO™**:

- in combination with copper containing fungicides, phosphates or phosphonates
- within 30 days after application of any copper fungicides
- in combination with highly alkaline material such as lime sulphur or certain boron products.

CALMAG-PRO™ should preferably be applied in the early morning or late afternoon. Do not apply to plants that are undergoing a period of moisture or heat stress.

For optimum results, **CALMAG-PRO™** should be used in a programme and in combination with other balanced fertilizers and foliar feeds. Consult your chemical distributor should crop specific programmes or any other information be required about the use of **CALMAG-PRO™**.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAX. RATE PER HA.	RATE PER 100ℓ WATER	REMARKS
Apples	11,5 ℓ	700 mℓ	Apply as a full cover spray in 1000 to 3000 ℓ spray water per hectare directly after fruit set. Repeat every 7 to 10 days to ensure 6 to 8 sprays and 6 to 12 sprays for bitter pit sensitive cultivars. If vigorous shoot growth appears use the higher dosage and increase the amount of sprays.
Avocado's	11,5 ℓ	700 mℓ	Apply as a full cover spray in 1000 to 3000 ℓ spray water per hectare. Start first application on flower trusses. Repeat 3 times in early spring on new growth. Add 6 mℓ Aqua-Wet per 100 ℓ water.
Cruciferae, Lettuce, Beet	4,5 ℓ		Apply 2 to 3 sprays, starting 4 weeks after transplant. Add Aqua-Wet at 6 mℓ per 100 ℓ. Apply in 300 ℓ water per ha.
Celery	4,5 ℓ		Start application after seedlings have established. Repeat every 7 to 10 days. Add Aqua-Wet at 6 mℓ per 100 ℓ. Apply in 300 ℓ water per hectare.
Cucurbits, Tomatoes & Peppers	6 ℓ		Start application after fruit set. Repeat every 7-10 days. Add Aqua-Wet at 6 mℓ per 100 ℓ. Apply in 500 ℓ water per ha.
Citrus	11,5 ℓ	700 mℓ	Start application in October directly after fruit set. Repeat every 7-10 days (4-5 applications). Apply in 1000 -3000 ℓ water per ha.
Cotton	4,5 ℓ		Start application when plants reach a height of 30 cm. Apply 2-3 sprays. Add Aqua-Wet at 6 mℓ per 100 ℓ. Apply in 500 ℓ water per hectare.
Grapes	6 - 8 ℓ		Start application directly after fruit set. Apply 3-4 sprays. Apply in 1000-1500 ℓ water per hectare.
Litchi & Mangoes	11,5 ℓ	700 mℓ	Start application during flowering. Apply 2-3 sprays at 10-day intervals. Add Aqua-Wet at 6 mℓ per 100 ℓ. Apply in 1000-3000 ℓ water per hectare.
Potatoes	6 ℓ		Start application at tuber initiation. Apply 3-4 sprays at 10-14 day intervals. Add Aqua-Wet at 6 mℓ per 100 ℓ. Apply in 500 ℓ water per hectare.
Soybeans, Groundnuts, Green Beans	4,5 ℓ		Start application at flowering. Apply 2 sprays at 7-day intervals. Add Aqua-Wet at 6 mℓ per 100 ℓ. Apply in 300 ℓ water per ha.
Stone Fruit (Plums, Peaches, Nectarines)	4,5 - 9 ℓ	350 mℓ	Start application directly after fruit set. Apply 4-6 sprays at 10-day intervals. Add Aqua-Wet at 6 mℓ per 100 ℓ. Apply in 1000-3000 ℓ water per hectare.
Sugarcane	6 ℓ		Start application in early spring. Apply 2 sprays at 10-day intervals. Add Aqua-Wet at 6 mℓ per 100 ℓ. Apply in 500-1500 ℓ water per hectare.
Other Crops	6 ℓ		Apply as a medium cover spray and repeat at 14-day intervals. Apply in 500 ℓ water per hectare.

Use lower rates as a maintenance spray and higher rates to correct deficiencies.