

# GLUCO-K™

**GLUCO-K™ is an organic potassium fertilizer for use as a foliar feed to maintain or correct potassium levels in plants**

## ACTIVE INGREDIENTS

Potassium (K)	184 g/kg	(230 g/ℓ)
L-amino acids	21 g/kg	( 26 g/ℓ)

**SG = 1,26 ±0,02**

*Gluconate Chelated*

**Sineria**

SINERIA HOLLAND BV  
Randwycksingel 20-A015  
6229 EE Maastricht The Netherlands  
[www.sineria.com](http://www.sineria.com)

Contents • Batch number • Manufacture date

Blank area for labeling information.

For optimum results, **GLUCO-K™** should be used in a programme and in combination with other balanced fertilizers and foliar feeds. **GLUCO-K™** is compatible with most agricultural remedies. It is however advisable to do a miscibility test prior to mixing with other chemicals. **Do not mix GLUCO-K™** with highly alkaline material such as Lime Sulphur and Bordeaux mixture, or with any phosphate-containing fertilizers. **GLUCO-K™** should preferably be applied in the early morning or late afternoon. Do not apply to plants undergoing a period of moisture or heat stress. Consult your chemical distributor should crop specific programmes or any other information be required about the use of **GLUCO-K™**.

## DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAX. RATE PER HA.	RATE PER 100ℓ WATER	REMARKS
Avocado, Mango and Litchi	10 ℓ	2 ℓ	1 to 2 applications on flower tissues and repeat on young summer growth. Apply in a minimum of 1000 - 2000 ℓ water per hectare.
Citrus	5 ℓ	0,5 - 1 ℓ	Two applications starting mid-November and repeat 21 days later. Apply as a medium cover spray in a minimum of 1000 - 3000 ℓ water per hectare.
Grapes	5 ℓ	1 - 2 ℓ	Apply after fruit set when fruit is pea-sized and repeat 21 days later in a minimum of 1000 ℓ water per hectare.
Grains, Peas & Canola	5 ℓ	0,5 - 1 ℓ	Four applications after flowering with 7 to 14 day intervals. Apply in a minimum of 500 ℓ water per hectare.
Pome and Stone Fruit	5 ℓ	1 ℓ	2 to 3 applications after fruit set, during fruit expansions, and repeat at 7 to 14 day intervals. Apply in a minimum of 1000 - 2000 ℓ water per hectare.
Potatoes	5 ℓ	0,5 - 1 ℓ	Apply after tuber swelling and repeat at 7 to 14 day intervals. Apply in a minimum of 500 ℓ water per hectare.
Tobacco	10 - 15 ℓ	2 ℓ	2 to 3 applications after topping in a minimum of 500 - 750 ℓ water per hectare.
Vegetables (Tomatoes, Eggplant, Cucurbits, Beet, Peppers, Lettuce, Carrots)	2,5 ℓ	250 - 500 ml	4 to 5 applications directly after fruit formation. Continue at 7 to 14 day intervals until 2 weeks prior to harvesting. Apply in a minimum of 500 ℓ water per hectare.

Use lower rates as a maintenance spray and higher rates to correct deficiencies.