

GERMINATOR COPPER-ZINC™

60.0% W.V. (600g/L) ZINC
15.0% W.V. (150g/L) COPPER
PLUS 10% GERMINATION HORMONES

**HIGH ANALYSIS ZINC AND
 COPPER SUSPENSION WITH
 GERMINATION BOOSTER, ESPECIALLY
 DESIGNED FOR SEED TREATMENT & EARLY
 FOLIAR APPLICATION IN ALL BROADACRE CROPS**

Supplier:

Sineria

SINERIA HOLLAND BV
 Randwycksingel 20-A015
 6229 EE Maastricht
 The Netherlands
www.sineria.com

Contents :	
Date of Manufacture:	
Date of Expiry:	
Batch No.	

READ ENTIRE LABELS AND ADDITIONAL WARRANTY SECTIONS BEFORE USE

GERMINATOR COPPER-ZINC™

ANALYSIS:

			W.V.%
Zinc	(Zn)	as Zinc Oxide	60.0
Copper	(Cu)	as Copper Oxide	15.0
Germination Booster (Ecklonia maxima)			10.0
	Added Water	69.4% W.V.	
	Total Solids	85.0% W.V.	

GERMINATOR Copper-Zinc™ is compatible with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity and observe precipitation (fall out) in mixture. If precipitation occurs, DO NOT proceed with application.

MIX ENOUGH GERMINATOR COPPER-ZINC™ FOR 1 DAY'S APPLICATION. DO NOT STORE GERMINATOR COPPER-ZINC™ MIXED WITH WATER OR OTHER CHEMICALS.

CONDITIONS OF SALE

1. All goods supplied are made from high grade materials and are believed to be suitable for use.
2. As no control can be exercised over storage, handling, mixing, application or use, or weather, plant or soil conditions before, during or after application (and all of which may affect the performance of the goods), NO RESPONSIBILITY for or liability for any failure in performance, losses, damages, or injuries (consequential or otherwise) arising from such storage, mixing, application or use, will be accepted under any circumstances whatsoever, and buyer assumes all responsibility for use of the product.
3. These conditions cannot be modified, varied or waived by our staff, distributors, or retailers, whether or not they advise or assist in the storage, handling, mixing or use of the goods, and such persons shall be entitled to the benefit of Clause 2.
4. MAXIMUM LIABILITY. Liability is limited to replacement of faulty goods only.

Net Contents	5 Litre	10 Litre	100 Litre	200 Litre	1000 Litre
--------------	---------	----------	-----------	-----------	------------

DIRECTIONS FOR USE:

AGITATE CONTENTS WELL BEFORE DILUTION

SUITABLE FOR APPLICATION BY: Boom spray and seed dressing units

SEED TREATMENT

CROP	RATE
WHEAT, BARLEY & COTTON	4-5L / TONNE SEED
MAIZE & RICE	5-8L / TONNE SEED
GRAIN & LEGUME	5-6L / TONNE SEED

Note: If using the lower rate, follow-up tissue tests may be required to determine the need for Foliar Application after emergence

FOLIAR TREATMENT

NOTE: Not compatible with inoculants

CROP	RATE
Broadhectare Crops	0.2 - 0.8 L/Ha

STANDARD MIXING & SPRAYING INSTRUCTIONS

1. Half fill the spray tank with clean water. Add the required quantity of GERMINATOR COPPER-ZINC™.
2. Fill the spray tank with water, as well as the amount of oil needed, to the required level while maintaining agitation to ensure thorough mixing of the spray mixture before spraying commences. Maintain agitation while spraying.
3. Tank mixtures must be sprayed out immediately and not allowed to stand in the spray tank overnight.
4. Ensure that agitation takes place for a few minutes prior to commencement of spraying after normal operational stops, e.g. lunch break.
5. Spraying equipment must be thoroughly flushed out at the end of the spraying operation.

COMPATIBILITY: Germinator Copper-Zinc™ is compatible with SEED PLUS & Vitavax Combination of the above seed coating products and Germinator Copper-Zinc™ is permissible.