

IRON-PRO™

IRON-PRO™ is an organic iron fertilizer for use as a foliar feed to maintain or correct iron levels in plants

FERTILIZER GROUP 2

ACTIVE INGREDIENT

Iron (Fe) 60 g/kg (74 g/l)

SG = 1,23 ± 0,02

L-amino acid chelated

Sineria

SINERIA HOLLAND BV
Randwycksingel 20-A015
6229 EE Maastricht The Netherlands
www.sineria.com

Contents • Batch number • Manufacture date

Empty box for labeling information.

IRON-PRO™ is compatible with most agricultural remedies. However, do **NOT** apply **IRON-PRO™**:

- in combination with copper containing fungicides, phosphates or phosphonates
- within 30 days after application of any copper fungicides
- in combination with highly alkaline material such as lime sulphur or certain boron products.

IRON-PRO™ should preferably be applied in the early morning or late afternoon. Do not apply to plants that are undergoing a period of moisture or heat stress.

For optimum results, **IRON-PRO™** should be used in a programme and in combination with other balanced fertilizers and foliar feeds. Consult your chemical distributor should crop specific programmes or any other information be required about the use of **IRON-PRO™**.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAX. RATE PER HA.	RATE PER 100ℓ WATER	REMARKS
Citrus	1,5 - 6 ℓ	80 - 130 mℓ	Two applications with 7 to 14 day intervals. Do not apply after fruit set. Apply in 2000 to 5000 ℓ water per hectare.
Field and Vegetable Crops	1,5 - 2,5 ℓ	550 mℓ	Apply during early growth stage when enough leaves have developed for absorption. Apply in 300 to 500 ℓ water per hectare.
Grapes	0,5 - 2 ℓ	80 - 130 mℓ	Apply weekly until deficiency symptoms disappear. Do not apply during flowering. Apply in 1000 to 1500 ℓ water per hectare.
Pome & Stone Fruit	1,5 - 6 ℓ	80 - 130 mℓ	Three applications with 7 to 10 day intervals. Apply first application when first leaves appear in spring. Apply in 2000 to 5000 ℓ water per hectare.
Sugar Cane	1 - 2 ℓ	350 mℓ	Apply in 300 to 500 ℓ water per hectare. Aerial application in 30 to 40 ℓ water per hectare. Apply early in the season when enough leaves have developed for absorption.

Use lower rates as a maintenance spray and higher rates to correct deficiencies.