

MAG-PRO™

MAG-PRO™ is an organic magnesium fertilizer for use as a foliar feed to maintain or correct magnesium levels in plants

FERTILIZER GROUP 1

ACTIVE INGREDIENT

Magnesium (Mg) 43 g/kg (56 g/ℓ)

SG = 1,32 ± 0,02

L-amino acid chelated

Sineria

SINERIA HOLLAND BV
Randwycksingel 20-A015
6229 EE Maastricht The Netherlands
www.sineria.com

Contents • Batch number • Manufacture date

--

MAG-PRO™ is compatible with most agricultural remedies. However, do **NOT** apply **MAG-PRO™** :

- in combination with copper containing fungicides, phosphates or phosphonates
- within 30 days after application of any copper fungicides
- in combination with highly alkaline material such as lime sulphur or certain boron products.

MAG-PRO™ should preferably be applied in the early morning or late afternoon. Do not apply to plants that are undergoing a period of moisture or heat stress.

For optimum results, **MAG-PRO™** should be used in a programme and in combination with other balanced fertilizers and foliar feeds. Consult your chemical distributor should crop specific programmes or any other information be required about the use of **MAG-PRO™**.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAX. RATE PER HA.	RATE PER 100ℓ WATER	REMARKS
Citrus	9 ℓ	350 - 700 mℓ	Apply 1 to 2 applications as a high volume spray in a minimum of 1000 ℓ water per hectare.
Cotton	4 ℓ	350 - 700 mℓ	Apply 1 to 2 applications as soon as deficiency symptoms appear. Apply in 300 - 500 ℓ water per ha.
Grapes	7 ℓ	350 - 700 mℓ	Start application two weeks after flowering. Repeat with 14 day intervals in a minimum of 1000 ℓ water per ha.
Grains, Potatoes, Peas, Canola	4 - 4,5 ℓ	350 - 700 mℓ	Start application when deficiency symptoms appear. Repeat if necessary. Apply in a minimum of 300 - 500 ℓ water per hectare.
Onions	4,5 ℓ	350 - 700 mℓ	Apply 3 to 4 applications. Apply in a minimum of 500 ℓ water per hectare.
Tea	8,5 ℓ	350 - 700 mℓ	Apply as a ground spray as soon as deficiency symptoms appears in 200 - 500 ℓ water per hectare. Aerial application in 30 - 45 ℓ water per hectare.
Tobacco	4 ℓ	350 - 700 mℓ	Apply 3 to 4 applications at 10 to 14 day intervals. Apply in 500 ℓ water per hectare.
Cabbage, Cucurbits, Eggfruit, Lettuce, Peppers & Tomatoes	4,5 - 7 ℓ	350 - 700 mℓ	Apply 3 to 4 applications. Start application when deficiency symptoms appear. Apply in a minimum of 200 - 400 ℓ water per hectare.
Other Crops	2 - 6 ℓ	350 - 700 mℓ	Start application when deficiency symptoms appear and repeat if necessary. Apply in 300 - 500 ℓ water per ha.

Use lower rates as a maintenance spray and higher rates to correct deficiencies.