

MAN-PRO™

MAN-PRO™ is an organic manganese fertilizer for use as a foliar feed to maintain or correct manganese levels in plants

FERTILIZER GROUP 2

ACTIVE INGREDIENT

Manganese (Mn) 80 g/kg (109 g/l)

SG = 1,24 ± 0,02

L-amino acid chelated

Sineria

SINERIA HOLLAND BV
Randwycksingel 20-A015
6229 EE Maastricht The Netherlands
www.sineria.com

Contents • Batch number • Manufacture date

--

MAN-PRO™ is compatible with most agricultural remedies. However, do **NOT** apply **MAN-PRO™**:

- in combination with copper containing fungicides, phosphates or phosphonates
- within 30 days after application of any copper fungicides
- in combination with highly alkaline material such as lime sulphur or certain boron products.

MAN-PRO™ should preferably be applied in the early morning or late afternoon. Do not apply to plants that are undergoing a period of moisture or heat stress.

For optimum results, **MAN-PRO™** should be used in a programme and in combination with other balanced fertilizers and foliar feeds. Consult your chemical distributor should crop specific programmes or any other information be required about the use of **MAN-PRO™**.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAX. RATE PER HA.	RATE PER 100ℓ WATER	REMARKS
Citrus	1,5 ℓ	50 - 70 mℓ	Apply on new flush in spring. Repeat if necessary. Apply in a minimum of 1000 - 3000 ℓ water per hectare.
Dry beans	2 - 2,5 ℓ	50 - 70 mℓ	Start application early in the growing season when sufficient leaves have developed for absorption. Repeat 14 days later if necessary. Apply in a minimum of 300 ℓ water per hectare.
<u>Field Crops</u> (Grain, Peas, Soybeans, Groundnuts, Crucifers, Cucumbers, Tomatoes, Chilies, Potatoes)	2,5 ℓ	50 - 70 mℓ	Start application early in the growing season when sufficient leaves have developed for absorption. Repeat three to four weeks later if necessary. Apply in a minimum of 300 ℓ water per hectare.
Olives	1,5 ℓ	50 - 70 mℓ	Apply as a foliar spray during late spring. Apply in a minimum of 1000 - 3000 ℓ water per hectare.
Pomefruit (Apples and Pears)	1,5 ℓ	50 - 70 mℓ	Apply as a foliar spray before blossom and repeat after harvest. Apply in a minimum of 1000 - 3000 ℓ water per hectare.
Pastures	5 ℓ	50 - 70 mℓ	Apply in early spring. Apply in a minimum of 300 ℓ water per hectare.
Stonefruit (Apricots, Peaches, Plums)	1,5 ℓ	50 - 70 mℓ	Apply as a foliar spray after harvest in a minimum of 1000 - 3000 ℓ water per hectare.

Use lower rates as a maintenance spray and higher rates to correct deficiencies.