

ZMC-PRO™ is compatible with most agricultural remedies. However, do **NOT** apply **ZMC-PRO™** :

- in combination with copper containing fungicides, phosphates or phosphonates
- within 30 days after application of any copper fungicides
- in combination with highly alkaline material such as lime sulphur or certain boron products.

ZMC-PRO™ should preferably be applied in the early morning or late afternoon. Do not apply to plants that are undergoing a period of moisture or heat stress.

For optimum results, **ZMC-PRO™** should be used in a programme and in combination with other balanced fertilizers and foliar feeds. Consult your chemical distributor should crop specific programmes or any other information be required about the use of **ZMC-PRO™**.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAX. RATE PER HA.	RATE PER 100ℓ WATER	REMARKS
Apples and Pears	3 ℓ	150 mℓ	Apply 2 to 3 applications, one early in the season and again after harvest in a minimum of 1000 to 3000 ℓ water per hectare.
Avocado's	3 ℓ	150 mℓ	Start first application on flowers in the spring and on all new growth in the summer. Repeat 2 to 3 times if necessary. Apply in 1000 to 3000 ℓ water per hectare.
Beans, Groundnuts, Peas, Soybeans	1,5 ℓ		Apply at flowering. Repeat twice at 7 to 10 day intervals if necessary. Apply in 300 ℓ water per hectare.
Cruciferae, Lettuce, Beetroot and Celery	1,5 ℓ		Start application 4 weeks after transplant and repeat 2 to 3 times if necessary. Apply in 300 to 500 ℓ water per hectare.
Citrus	3 ℓ	150 mℓ	Apply as a full cover spray in spring to all new growth and repeat 2 to 3 times. Do not spray directly before or during harvest. Apply in 1000 to 3000 ℓ water per hectare.
Cotton	2 ℓ		Apply early in the season at plant height of 30 cm, and repeat 2 to 3 times. Apply in 500 ℓ water per hectare.
Cucurbits (Pumpkins, etc.)	1,5 ℓ		Start application early in spring, and repeat every 7 to 10 days if necessary. Apply in 500 ℓ water per hectare.
Macadamia, Pecans and other Nuts	3 ℓ	100 mℓ	Do first application in the spring and then on all new growth later in the season or when deficiencies occur. Two to three applications are recommended. Apply in 4000 ℓ water per hectare.
Maize and Sweet Corn	1,5 ℓ	50 mℓ	Do first application early in the season at the first signs of a deficiency and repeat the application if required. Apply in 200 ℓ water per hectare.
Mangoes	3 ℓ	150 mℓ	Start application shortly before flowering. Apply 2 to 3 sprays at 10 day intervals. Apply in 1000 to 3000 ℓ water per hectare.
Olives	1,5 ℓ	75 mℓ	Apply early in the growing season and repeat 3 to 4 weeks later on new growth. Apply in 500 ℓ water per hectare.
Onions	1,5 ℓ	75 mℓ	Apply at the first signs of a deficiency. Repeat two weeks later if required. Apply in 500 ℓ water per hectare.
Potatoes, Tomatoes, Peppers, Chillies	1,5 ℓ	75 mℓ	Apply at the first signs of a deficiency and repeat 3 to 4 weeks later if necessary. Apply in 500 ℓ water per hectare.
Stone Fruit	1,5 ℓ	75 mℓ	Apply very early in the season before blossom and then again after harvest. Apply in 500 ℓ water per hectare.
Tea	1 ℓ	45 mℓ	3 to 4 applications depending on the severity of the deficiency on new spring growth. Apply in 1000 to 2000 ℓ water per hectare.
Wheat, Barley, Oats	2 - 3 ℓ	150 mℓ	Apply early in the season at the first signs of a deficiency and repeat the application if required. Apply in 200 ℓ water per hectare.
Other Crops	2 ℓ	100 mℓ	Apply as a full cover spray and repeat every 14 days as required. Apply in 500 ℓ water per hectare.

Use lower rates as a maintenance spray and higher rates to correct deficiencies.