

ZINC-PRO™

ZINC-PRO™ is an organic zinc fertilizer for use as a foliar feed to maintain or correct zinc levels in plants

FERTILIZER GROUP 2

ACTIVE INGREDIENT

Zinc (Zn) 100 g/kg (142 g/l)

SG = 1,40 ± 0,02

L-amino acid chelated

Sineria

SINERIA HOLLAND BV
Randwycksingel 20-A015
6229 EE Maastricht The Netherlands
www.sineria.com

Contents • Batch number • Manufacture date

--

ZINC-PRO™ is compatible with most agricultural remedies. However, do **NOT** apply **ZINC-PRO™**:

- in combination with copper containing fungicides, phosphates or phosphonates
- within 30 days after application of any copper fungicides
- in combination with highly alkaline material such as lime sulphur or certain boron products.

ZINC-PRO™ should preferably be applied in the early morning or late afternoon. Do not apply to plants that are undergoing a period of moisture or heat stress.

For optimum results, **ZINC-PRO™** should be used in a programme and in combination with other balanced fertilizers and foliar feeds. Consult your chemical distributor should crop specific programmes or any other information be required about the use of **ZINC-PRO™**.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	MAX. RATE PER HA.	RATE PER 100€ WATER	REMARKS
Apples & Pears	3 ℓ	45–90 mℓ	Two applications, one early in the season and again after harvest in 500 ℓ to 2000 ℓ water per ha.
Avocado's	7 ℓ	70–135 mℓ	Start first application on flowers in the spring and on all new growth in the summer. Two to three applications are recommended. Apply in 2000 ℓ water per hectare.
Beans, Groundnuts, Peas, Soybeans	2 ℓ	45–90 mℓ	One to two applications early in the growing season. Apply in 200 ℓ water per hectare.
Cruciferae (Cabbages, etc.)	2 ℓ	45–90 mℓ	Apply at the first signs of a deficiency and repeat 3 to 4 weeks later if necessary. Apply in 500 ℓ water per ha.
Citrus	3 ℓ	45–90 mℓ	Apply two to three applications as a full cover spray in spring to all new growth. Do not spray directly before or during harvest. Apply in 2000 ℓ water per hectare.
Cotton	2 ℓ	45–90 mℓ	Do first application early in the season and repeat the application if required. Apply in 500 ℓ water per hectare.
Cucurbits (Pumpkins, etc.)	2 ℓ	45–90 mℓ	Apply at the first signs of a deficiency and repeat 3 to 4 weeks later. Apply in 500 ℓ water per hectare.
Lettuce	2 ℓ	45–90 mℓ	One to two applications early in the growing season. Apply in 500 ℓ water per hectare.
Macadamia, Pecans and other Nuts	7 ℓ	70–135 mℓ	Apply the first application in the spring and then on all new growth later in the season or when deficiencies occur. Two to three applications are recommended. Apply in 4000 ℓ water per ha.
Maize, Sweet Corn and Small Grains	2 ℓ	45–90 mℓ	Apply first application early in the season at the first signs of a deficiency and repeat the application if required. Apply in 200 ℓ water per hectare.
Mangoes	3 ℓ	70–135 mℓ	Apply as a full cover spray in the spring and repeat the application in the summer if required. Apply in 3000 to 4000 ℓ water per ha.
Olives	2 ℓ	90 mℓ	Apply early in the growing season and repeat 3 to 4 weeks later on new growth. Apply in 500 ℓ water per ha.
Onions	2 ℓ	45–90 mℓ	Apply at the first signs of a deficiency. Repeat two weeks later if required. Apply in 500 ℓ water per ha.
Solanaceae (Peppers, etc.)	2 ℓ	45–90 mℓ	Apply at the first signs of a deficiency and repeat 3 to 4 weeks later if necessary. Apply in 500 ℓ water per ha.
Stone Fruit (Peaches, etc.)	1,5 ℓ	70–135 mℓ	Apply very early in the season and then again after harvest. Apply in 500 ℓ water per ha.
Tea	2 ℓ	45–90 mℓ	Three to 4 applications depending on the severity of the deficiency on new spring growth. Apply in 1000 to 2000 ℓ water per ha.
Other Crops	2 ℓ	45–90 mℓ	Apply as a full cover spray and repeat every 14 days as required. Apply in 500 ℓ water per hectare.

Use lower rates as a maintenance spray and higher rates to correct deficiencies.