# Agri Star 75WDG<sup>TM</sup>

Agristar 75WDG™ is a broad-spectrum, post-emergence herbicide with some soil residual activity for broadleaf weed control in wheat.

#### Agri-Star 75WDG™ advantages

- High active ingredient content in a special granular formulation offering convenient storage, transport and application.
- Wide application window
- High safety in application
- Reliable and effective control of a wide range of broadleaf weeds
- Excellent crop tolerance
- Excellent crop rotation options
- WHO Classification: U unlikely to present acute hazard in normal use

## Sineria

### Agri-Star 75WDG™ Mode of Action

- Agristar 75WDG™ is rapidly absorbed by the plant foliage and roots and is translocated throughout the plant. Susceptible weeds cease growth almost immediately after post-emergence treatment and are killed in 7-21 days.
- Warm, moist conditions following treatment promote the activity of Agristar 75WDG™.
- Sulfonylurea herbicide acting as an ALS inhibitor
- WSSA Group: 2

Crop	Diseases	Rate per Ha
Wheat	Broadleaf weeds	15 gr/ha or 12.5 gr/ha if tank mixed

Application Timing: Apply from the 3-leaf stage of the wheat up to the fully emerged flag leaf stage.

#### **Important Notes**

- Agristar 75 WDG<sup>™</sup> should not be used where the following weeds are dominant: Yellow daisy (*Chrysanthemum* sp.) and Cleavers (*Gallium* sp.).
- Do not apply Agristar 75 WDG<sup>™</sup> when stressed by serious weather conditions, drought, disease or insect damage as crop injury may result.

#### **Active Ingredient**

Tribenuron-methyl 750 gr/kg

#### **Common Name** Scientific Name Oxalis Oxalis sp. **Spurrey** Spergula arvensis Chickweed Stellaria media Wild radish Raphanus raphanistrum **Black bindweed** Fallopia convolvulus Goosefoot Chenopodiumsp. Knotweed Convolvulussp. Black nightshade Solanum nigrum **Blackjack Weed** Bidens pilosa Double thorn Oxygonum sinuatum

#### **Recommended Application Rates**

- ► Half-fill spray tank with water. Premix the required amount of Agristar 75WDG<sup>TM</sup> with a small amount of water. Add this premix to the tank. Top up tank with water to the required volume, then agitate and mix thoroughly.
- Water volumes of 100-200 liter per hectare for boom application and 30-50 liter for aerial application are recommended
- For the best results spray when the broadleaf weeds are relatively small and vigorously growing.

Common Name	Scientific Name
Shepherd's purse	Capsella bursa-pastoris
Pigweed	Amaranthussp.
Mallow	Malva sp.
Apple of Peru	Nicandra physalodes
Mexican marigold	Tagetes sp.
Gallant soldier	Galinsoga parviflora
Devil's thorn	Tribulus terrestris
Jimson weed (Thorn Apple)	Datura stramonium
Common purslane	Portulaca oleracea
Mayweed	Anthemissp.

Disclaimer: This information and all further technical advice is based on our present knowledge and experience and approvable from the registration authorities. However, it implies no liability or other legal responsibility on our part acluding with regard to existing third party intellectual property rights, especially patent rights, or particular, no warranty whether express or implied, or guarantee of product properties in the legal sense is intended or implied. We reserve the light to make any changes according to technological progress or further developments, in the event of an discrepancies between the information stated herein or any other information source and the information stated on the able of the product, the information stated on the heads of the product will prevail. The customer is not released from the obligation to conduct careful inspection and testing of incoming goods. Performance of the product describes terein should be verified by testing, which should be carried out only by qualified experts in the sole responsibility of the sustance. Reference to trade names use by other companies is neither a recommendation, nor does it imply that similar products could not be used.





SINERIA Holland BV Randwycksingel 20-A015 6229 EE Maastricht, The Netherlands