




Embark 90SC

Flonicamid:

- Pyridinocarboxamide insecticide

- N-(cyanomethyl)-4-(trifluoromethyl)pyridine-3-carboxamide




- Chemical Formula: $C_9H_6F_3N_3O$

Flonicamid is a systemic insecticide with a novel mode of action.

Flonicamid's mode of action is different from any other insecticide. Flonicamid belongs to IRAC Group 29 and is the only insecticide in this class.

Emamectin-benzoate:

- Avermectin insecticide
- 4''-Deoxy-4''-epi-methylaminoavermectin B1; Epimethylamino-4''-deoxyavermectin



Emamectin B_{1a}
R = CH₂CH₃

Emamectin B_{1b}
R = CH₃

- Chemical Formula: $C_{49}H_{75}NO_{13}$

Emamectin-benzoate is a natural fermentation product obtained from soil microorganisms. Emamectin-benzoate has a unique mode of action which is different from any other insecticide. Acts by interfering with the pest's nervous system, causing them to lose cell function and stop feeding. The larvae die in 2-4 days.
IRAC Group: 6 (Chloride channel activators)

Soft chemistry, novel, contact and stomach insecticide, with systemic and translaminar activity for the control of insect pests and spidermites on a wide range of crops.

- Unique synergetic mixture of a Synthetic and a Fermentation product
- Contact, translaminar and systemic action

IPM Compatible

- Embark 90SC has relatively low impact on beneficial organisms following application.
- Always apply late in the evening when bees are not foraging.
- Embark 90SC has a good toxicological, environmental and eco-toxicological profile
- Flonicamid exhibits excellent performance for control of almost all important soft body insects
- Flonicamid has no negative impact on pollinating insects or natural enemies.
- Flonicamid provides a new option for integrated pest management programs due to its soft toxicological, and eco-toxicological profile.

Application timing

- Embark 90SC is recommended to be applied upon the pest appearance
- Ensure thorough coverage of the canopy with the spraying mixture.

In case of high pest pressure, repeat application at 7-10 days intervals, with no more than 4 applications per growing season.

Directions For Use:

Embark 90SC should be mixed with water to be ready for use. It may be applied with any type of spraying equipment.

Rate of Application

CROPS	PEST	RATE
Ornamentals	Aphids, Thrips, Whiteflies, Mealybugs, Leafhoppers, Spidermites, Leafminers, Lepidoptera pests	0.25 - 0.35 Lt/Ha
Vegetables (Peppers, Tomato, Cruciferae, Carrots, Paprika, Peppers, Eggplant)	Aphids, Thrips, Whiteflies, Mealybugs, Leafhoppers, Spidermites, Leafminers, Lepidoptera pests	0.25 - 0.35 Lt/Ha
Potatoes, Beans, Soybean	Aphids, Thrips, Whiteflies, Mealybugs, Leafhoppers, Spidermites, Leafminers, Lepidoptera pests	0.25 - 0.35 Lt/Ha
Maize, Wheat	Aphids, Thrips, Whiteflies, Mealybugs, Leafhoppers, Spidermites, Leafminers, Lepidoptera pests	0.25 - 0.35 Lt/Ha

Always alternate with insecticides with different mode of action through the season to prevent the development of resistant pests.

Important Note:

The Indicated Crops and Recommended Rate of application on this Product informative sheet may not be applicable in country where the product is intended to be used. User must refer and use the product only as per the officially Register and approved uses by the ministry of Agriculture authorized authorities. The supplier cannot be held responsible or liable if the use of this product is performed on Crops which are not listed on the official label as approved by the ministry of Agriculture authorized authorities.

Disclaimer: This information and all further technical advice is based on our present knowledge and experience and approvals from the registration authorities. However, it implies no liability or other legal responsibility on part, including with regard to existing third party intellectual property rights, especially patent rights. In particular, no warranty, whether express or implied, or guarantee of product properties in the legal sense is intended or implied. We reserve the right to make any changes according to technological progress or further developments. In the event of any discrepancies between the information stated herein or any other information source and the information stated on the label of the product, the information stated on the label of the product will prevail. The customer is not released from the obligation to conduct careful inspection and testing of incoming goods. Performance of the product described herein should be verified by testing, which should be carried out only by qualified experts in the sole responsibility of the customer. Reference to trade names use by other companies is neither a nor does it imply that similar products could not be used. Embark 90 SC is a trademark from Sineria Kenya Ltd.

Embark 90 SC is a trademark from Sineria Kenya Ltd.

Field Trials

Extensive field research has proven the efficacy of Embark 90SC. In Hasselt, Belgium (2018) the following results were obtained:




Figure:
% Efficacy of Embark 90SC:
1 application at 250 ml/ha

