

FRUIT



- Category: Crop Nutrition Range
- Formulation type: Gel
- Guaranteed Analysis (w/v):
- Nitrogen (N) 8.0 %
 Phosphorus (P) 8.0 %
 Potassium (K) 60.0 %
 Boron (B) 0.002 %
 Iron* (Fe) 0.017%
 Copper* (Cu) 0.05%
 Manganese* (Mn) 0.005 %
- Zinc* (Zn) 0.005 % - Molybdenum (Mo) 0.0001 % * EDTA chelated
- Physical Characteristics:
- Density (at 20°C) 1.5 1.7
- Physical state Gel

NEW BREAKTHROUGH TECHNOLOGY From The Netherlands

08-08-60+ME

NutriGel FRUIT is a water soluble gel fertilizer with chelated micro-elements, specifically designed for use as foliar feeding or fertigation.

Features & Benefits:

- New generation water soluble Gel Fertilizer.
- Control release technology in patented Gel Matrix form.
- Positively charged Gel Matrix formulation that allows better lodging of the entire Nutrient on all negatively charged crops.
- NutriGel is a Gel Matrix that bonds to the leaf surface for a prolonged period of time.
- Effective translocation and metabolism.
- Compatible with most agrochemicals.

Patented technology and compatibility:

- **NutriGel** line of products are High analyses suspended particles in a special **Gel matrix** that deliver slow release uniformity.
- NutriGel SHOOT is compatible with most agricultural remedies.
 It is however advisable to do a miscibility test prior to mixing
 NutriGel SHOOT with other agricultural products.
- NutriGel SHOOT has a neutral pH.

Application Methods:

NutriGel SHOOT is designed for use as a foliar feed or fertigation and can be applied by:

Boom sprayer | Knapsack | Irrigation system



DIRECTIONS FOR USE (USE ONLY AS DIRECTED)

CROP	RATE/HA	REMARKS
Field Crops: Canola, Wheat, Barley, Maize, Sunflower, Soybeans, Sorghum, Groundnut, Cotton, Tobacco, Lucerne, Pastures, etc.	1 - 2 L	Apply as a full cover spray during the reproductive growth stage or when the requirement for Potassium is high. Repeat application 14 days later or as required. Apply in 200-300Lt water/ha
Vegetable Crops: Lettuce, Beans, Brassica, Cucurbits, Garlic, Onions, Peppers, Peas, Tomato, Celery, Beetroot, Carrot, Strawberries, etc.	1-2L	Apply as a full cover spray during the reproductive growth stage or when the requirement for Potassium is high. Alternate with IntroGel SET. Repeat application 14 days later or as required. Apply in 200-400Lt water/ha
Pome and Stone Fruits: Apples, Pears, Nectarines, Peaches, Plums, Apricots, etc.	2 - 3 L	Apply as a full cover spray during the reproductive growth stage or when the requirement for Potassium is high. Alternate with IntroGel SET. Repeat application 14 days later or as required. Apply in 2000-3000Lt water/ha
Subtropical Fruits: Citrus, Avocados, Litchis, Mangoes, Bananas, Pineapples, etc.	2 - 3 L	Apply as a full cover spray during the reproductive growth stage or when the requirement for Potassium is high. Alternate with IntroGel SET. Repeat application 14 days later or as required. Apply in 2000-3000Lt water/ha
Nut Trees: Almond, Macadamia, Pecan, Walnut, etc.	3 - 4 L	Apply as a full cover spray during the reproductive growth stage or when the requirement for Potassium is high. Alternate with IntroGel SET. Repeat application 14 days later or as required. Apply in 2000-3000Lt water/ha
Other: Olives, Grapes, Figs, Pomegranates, Berries, etc.	2 - 3 L	Apply as a full cover spray during the reproductive growth stage or when the requirement for Potassium is high. Alternate with IntroGel SET. Repeat application 14 days later or as required. Apply in 500-1500Lt water/ha

Storage:

- Store in the original, well-sealed container away from direct sunlight in a cool dry environment.
- · Keep away from children, uninformed people, animals and foodstuffs.
- Do not re-use the container.


Warranty:

Although this product has been extensively tested under large variety of conditions, the supplier does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic conditions as well as method, time and accuracy of application.

Disclaimer: This information and all further technical advice is based on our present knowledge and experience and approvals from the registration authorities. However, it implies no liability or other legal responsibility on our part, including with regard to existing third party intellectual property rights, especially patent rights. In particular, no warranty, whether express or implied, or guarantee of product properties in the legal sense is intended or implied. We reserve the right to make any changes according to technological progress or further developments. In the event of any discrepancies between the information stated herein or any other information source and the information stated on the label of the product, the information stated on the label of the product will prevail. The customer is not released from the obligation to conduct careful inspection and testing of incoming goods. Performance of the product described herein should be verified by testing, which should be carried out only by qualified experts in the sole responsibility of the customer. Reference to trade names use by other companies is neither a recommendation, nor does it imply that similar products could not be used.

Application Instructions

- NutriGel FRUIT is recommended for use during plant growth periods that require relatively high Potassium levels such as during fruit formation and fruit growth.
- NutriGel FRUIT should preferably be applied in the early morning or late afternoon.
- Do not apply to plants that are undergoing a period of moisture or heat stress, or during cold wet conditions.
- Do not apply on wet fruit.
- For optimum results, NutriGel FRUIT should be used in a balanced crop nutrition program.



- 1 High concentration Gel
- Uniform distribution on the sub-surfaces of the leaf
- Positively charge formulation allows better lodging on all negatively charged crops
- (5) Control release technology in a GEL form

NutriGel is a trademark from Sineria Holland BV.



